

T violation in $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay beyond standard model

T. M. Aliev^{a *}, A. Özpineci^{b †}, M. Savcı^{a ‡}, C. Yüce^a

^a Physics Department, Middle East Technical University, 06531 Ankara, Turkey

^b The Abdus Salam International Center for Theoretical Physics, I-34100, Trieste, Italy

Abstract

Using the most general, model independent form of effective Hamiltonian, the transverse polarization P_T of Λ in $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay is studied. It is observed that the averaged $\langle P_T \rangle$ is very sensitive to the existence of new physics and can attain large values, which can be measured at future colliders.

PACS numbers: 12.60.-i, 13.30.-a, 14.20.Mr

*e-mail: taliev@metu.edu.tr

†e-mail: ozpineci@ictp.trieste.it

‡e-mail: savci@metu.edu.tr

1 Introduction

Recently time-reversal (T) violation has been measured in the K^0 system [1]. However, the origin of T violation, as well as CP violation which also has been obtained experimentally in K^0 meson system, remains unclear. In the standard model (SM) both violations comes from a phase of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [2]. It is well known that SM predicts sizeable CP violation in the B^0 meson system (see [3]). Detecting CP violation constitutes one of the main research directions of the working B meson factories [4]. These machines have already provided the first evidence for CP violation in B system, namely $\sin 2\beta = 0.78 \pm 0.08$ [5]. In search of other sources of CP violation one needs to investigate new processes. In this work we study the T-violating effects in the baryonic $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ ($\ell = \mu, \tau$) decays using the most general form of the effective Hamiltonian. Large number of Λ_b baryons are expected to be produced in Tevatron, LHC, etc., which is a great opportunity to examine the SM predictions for various decay modes of Λ_b . These decays are also very sensitive to the new physics beyond the SM. The interest to baryonic decays can be attributed to the fact that only these decays can give valuable information about the handedness of the quarks. It should be noted that T-violating effects in the $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ and $B \rightarrow K^* \ell^+ \ell^-$ decays were studied in the Supersymmetric model in [6] and [7], respectively.

It is well known that for a general three-body decay the triple spin-momentum correlations $\vec{s} \cdot (\vec{p}_i \times \vec{p}_j)$ are T-odd observables, where $\vec{s}$, $\vec{p}_i$ and $\vec{p}_j$ are the spin and momenta of final particles. So, in order to organize T-odd quantities we have two possibilities:

- either we choose lepton polarization as the polarization of final particles, or
- we choose only Λ baryon polarization.

In the $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay, experimentally only one polarization state of the particles can be measured. In principle one can use either polarization of the lepton or polarization of the baryon in studying T-odd observables. The first possibility, i.e., the lepton polarizations in the $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay was studied in detail in [8]. Therefore, in the present work we prefer choose the second possibility, namely, Λ baryon polarization in investigating the T-violating effects. It should be noted here that, lepton polarization effects for the $B \rightarrow K^* \ell^+ \ell^-$ were studied comprehensively in [9].

The paper is organized as follows. In section 2, using the most general, model independent form of the effective Hamiltonian, we derive the expression for the T-odd transverse polarization of Λ . Section 3 is devoted to the numerical analysis and concluding remarks.

2 Calculation of the transversal polarization of Λ baryon

The matrix element of the $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay at quark level is described by the $b \rightarrow s \ell^+ \ell^-$ transition. The decay amplitude for the $b \rightarrow s \ell^+ \ell^-$ transition, in a general model independent form, can be written in the following way (see [10, 11])

$$\mathcal{M} = \frac{G\alpha}{\sqrt{2}\pi} V_{tb} V_{ts}^* \left\{ C_{SL} \bar{s} i \sigma_{\mu\nu} \frac{q^\nu}{q^2} L b \bar{\ell} \gamma^\mu \ell + C_{BR} \bar{s} i \sigma_{\mu\nu} \frac{q^\nu}{q^2} b \bar{\ell} \gamma^\mu \ell + C_{LL}^{tot} \bar{s}_L \gamma^\mu b_L \bar{\ell}_L \gamma_\mu \ell_L \right.$$

$$\begin{aligned}
& + C_{LR}^{tot} \bar{s}_L \gamma^\mu b_L \bar{\ell}_R \gamma_\mu \ell_R + C_{RL} \bar{s}_R \gamma^\mu b_R \bar{\ell}_L \gamma_\mu \ell_L + C_{RR} \bar{s}_R \gamma^\mu b_R \bar{\ell}_R \gamma_\mu \ell_R \\
& + C_{LRLR} \bar{s}_L b_R \bar{\ell}_L \ell_R + C_{RLLR} \bar{s}_R b_L \bar{\ell}_L \ell_R + C_{LRRL} \bar{s}_L b_R \bar{\ell}_R \ell_L + C_{RLRL} \bar{s}_R b_L \bar{\ell}_R \ell_L \\
& + C_T \bar{s} \sigma^{\mu\nu} b \bar{\ell} \sigma_{\mu\nu} \ell + i C_{TE} \epsilon^{\mu\nu\alpha\beta} \bar{s} \sigma_{\mu\nu} s \sigma_{\alpha\beta} \ell \Big\} , \tag{1}
\end{aligned}$$

where $L = (1 - \gamma_5)/2$ and $R = (1 + \gamma_5)/2$ are the chiral operators and C_X are the coefficients of the four-Fermi interaction. Part of these Wilson coefficients and structures does already exist in the effective Hamiltonian of the $b \rightarrow s$ transition in the SM. The first two of the coefficients C_{SL} and C_{BR} are the nonlocal Fermi interactions which correspond to $-2m_s C_7^{eff}$ and $-2m_b C_7^{eff}$ in the SM, respectively. The following four terms describe vector type interactions. Two of these vector interactions containing the coefficients C_{LL}^{tot} and C_{LR}^{tot} do also exist in the SM in the forms $(C_9^{eff} - C_{10})$ and $(C_9^{eff} + C_{10})$, respectively. Therefore C_{LL}^{tot} and C_{LR}^{tot} represent the sum of the combinations from SM and the new physics in the following forms

$$\begin{aligned}
C_{LL}^{tot} &= C_9^{eff} - C_{10} + C_{LL} , \\
C_{LR}^{tot} &= C_9^{eff} + C_{10} + C_{LR} . \tag{2}
\end{aligned}$$

The terms with C_{LRRL} , C_{LRLR} , C_{RLRL} and C_{RLLR} describe the scalar type interactions. The last two terms in Eq. (1) correspond to the tensor type interactions. The amplitude of the exclusive $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay can be obtained by sandwiching the matrix element of the $b \rightarrow s \ell^+ \ell^-$ decay between initial and final state baryons. It follows from Eq. (1) that, in order to calculate the amplitude of the $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay the following matrix elements are needed

$$\begin{aligned}
& \langle \Lambda | \bar{s} \gamma_\mu (1 \mp \gamma_5) b | \Lambda_b \rangle , \\
& \langle \Lambda | \bar{s} \sigma_{\mu\nu} (1 \mp \gamma_5) b | \Lambda_b \rangle , \\
& \langle \Lambda | \bar{s} (1 \mp \gamma_5) b | \Lambda_b \rangle . \tag{3}
\end{aligned}$$

Explicit forms of these matrix elements in terms of the form factors are presented in Appendix-A of [12]. Using the parametrization of these matrix elements, the matrix form of the $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay can be written as

$$\begin{aligned}
\mathcal{M} &= \frac{G\alpha}{4\sqrt{2}\pi} V_{tb} V_{ts}^* \Big\{ \bar{\ell} \gamma^\mu \ell \bar{u}_\Lambda \Big[A_1 \gamma_\mu (1 + \gamma_5) + B_1 \gamma_\mu (1 - \gamma_5) \\
& + i \sigma_{\mu\nu} q^\nu [A_2 (1 + \gamma_5) + B_2 (1 - \gamma_5)] + q_\mu [A_3 (1 + \gamma_5) + B_3 (1 - \gamma_5)] \Big] u_{\Lambda_b} \\
& + \bar{\ell} \gamma^\mu \gamma_5 \ell \bar{u}_\Lambda \Big[D_1 \gamma_\mu (1 + \gamma_5) + E_1 \gamma_\mu (1 - \gamma_5) + i \sigma_{\mu\nu} q^\nu [D_2 (1 + \gamma_5) + E_2 (1 - \gamma_5)] \\
& + q_\mu [D_3 (1 + \gamma_5) + E_3 (1 - \gamma_5)] \Big] u_{\Lambda_b} + \bar{\ell} \ell \bar{u}_\Lambda (N_1 + H_1 \gamma_5) u_{\Lambda_b} + \bar{\ell} \gamma_5 \ell \bar{u}_\Lambda (N_2 + H_2 \gamma_5) u_{\Lambda_b} \\
& + 4 C_T \bar{\ell} \sigma^{\mu\nu} \ell \bar{u}_\Lambda \Big[f_T \sigma_{\mu\nu} - i f_T^V (q_\nu \gamma_\mu - q_\mu \gamma_\nu) - i f_T^S (P_\mu q_\nu - P_\nu q_\mu) \Big] u_{\Lambda_b} \\
& + 4 C_{TE} \epsilon^{\mu\nu\alpha\beta} \bar{\ell} \sigma_{\alpha\beta} \ell i \bar{u}_\Lambda \Big[f_T \sigma_{\mu\nu} - i f_T^V (q_\nu \gamma_\mu - q_\mu \gamma_\nu) - i f_T^S (P_\mu q_\nu - P_\nu q_\mu) \Big] u_{\Lambda_b} \Big\} , \tag{4}
\end{aligned}$$

where $P = p_{\Lambda_b} + p_\Lambda$.

Explicit expressions of the functions A_i , B_i , D_i , E_i , H_j and N_j ($i = 1, 2, 3$ and $j = 1, 2$) are given in Appendix-A of [12].

Obviously, the $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay introduces a lot of form factors. However, when the heavy quark effective theory (HQET) has been used, the heavy quark symmetry reduces the number of independent form factors to two only (F_1 and F_2), irrelevant with the Dirac structure of the relevant operators [13], and hence we obtain that

$$\langle \Lambda(p_\Lambda) | \bar{s} \Gamma b | \Lambda(p_{\Lambda_b}) \rangle = \bar{u}_\Lambda [F_1(q^2) + \not{q} F_2(q^2)] \Gamma u_{\Lambda_b} , \quad (5)$$

where Γ is an arbitrary Dirac structure, $v^\mu = p_{\Lambda_b}^\mu / m_{\Lambda_b}$ is the four-velocity of Λ_b , and $q = p_{\Lambda_b} - p_\Lambda$ is the momentum transfer. Comparing the general form of the form factors with (5), one can easily obtain the following relations among them (see also [14])

$$\begin{aligned} g_1 &= f_1 = f_2^T = g_2^T = F_1 + \sqrt{r} F_2 , \\ g_2 &= f_2 = g_3 = f_3 = g_T^V = f_T^V = \frac{F_2}{m_{\Lambda_b}} , \\ g_T^S &= f_T^S = 0 , \\ g_1^T &= f_1^T = \frac{F_2}{m_{\Lambda_b}} q^2 , \\ g_3^T &= \frac{F_2}{m_{\Lambda_b}} (m_{\Lambda_b} + m_\Lambda) , \\ f_3^T &= -\frac{F_2}{m_{\Lambda_b}} (m_{\Lambda_b} - m_\Lambda) , \end{aligned} \quad (6)$$

where $r = m_\Lambda^2 / m_{\Lambda_b}^2$. These relations will be used in further numerical calculations.

As has already been stated, in order to study the T-violating effects, we need spin polarization of Λ baryon. Spin vector s^μ of Λ baryon can be expressed in terms of the unit vector $\vec{\xi}$ along the Λ spin in its rest frame as,

$$s_\mu = \left(\frac{\vec{p}_\Lambda \cdot \vec{\xi}}{m_\Lambda}, \vec{\xi} + \frac{\vec{p}_\Lambda (\vec{p}_\Lambda \cdot \vec{\xi})}{m_\Lambda (E_\Lambda + m_\Lambda)} \right) , \quad (7)$$

and choose the unit vectors along the longitudinal, transversal and normal components of the Λ polarization to be

$$\begin{aligned} \vec{e}_L &= \frac{\vec{p}_\Lambda}{|\vec{p}_\Lambda|} , \\ \vec{e}_T &= \frac{\vec{p}_{\ell^-} \times \vec{p}_\Lambda}{|\vec{p}_{\ell^-} \times \vec{p}_\Lambda|} , \\ \vec{e}_N &= \vec{e}_L \times \vec{e}_T , \end{aligned} \quad (8)$$

respectively, where $\vec{p}_{\ell^-}$ and $\vec{p}_\Lambda$ are the three momenta of ℓ^- and Λ in the center of mass frame of the final leptons. The differential decay rate of the $\Lambda_b \rightarrow \Lambda \ell^+ \ell^-$ decay for any spin direction $\vec{\xi}$ along Λ baryon can be written in the following form

$$\frac{d\Gamma}{ds} = \frac{1}{2} \left(\frac{d\Gamma}{ds} \right)_0 \left[1 + \left(P_L \vec{e}_L + P_N \vec{e}_N + P_T \vec{e}_T \right) \cdot \vec{\xi} \right] , \quad (9)$$

where $(d\Gamma/ds)_0$ corresponds to the unpolarized differential decay rate, $s = q^2/m_{\Lambda_b}^2$ and P_L , P_N and P_T represent the longitudinal, normal and transversal polarizations of Λ , respectively. The unpolarized decay width in Eq. (9) can be written as

$$\left(\frac{d\Gamma}{ds}\right)_0 = \frac{G^2\alpha^2}{8192\pi^5} |V_{tb}V_{ts}^*|^2 \lambda^{1/2}(1, r, s) v \left[\mathcal{T}_0(s) + \frac{1}{3}\mathcal{T}_2(s)\right], \quad (10)$$

where $\lambda(1, r, s) = 1 + r^2 + s^2 - 2r - 2s - 2rs$ is the triangle function and $v = \sqrt{1 - 4m_\ell^2/q^2}$ is the lepton velocity. The explicit expressions for $\mathcal{T}_0$ and $\mathcal{T}_2$ can be found in [12].

The T-odd transverse polarization of Λ_b baryon is defined as

$$P_T(s) = \frac{\frac{d\Gamma}{ds}(\vec{\xi} = \vec{e}) - \frac{d\Gamma}{ds}(\vec{\xi} = -\vec{e})}{\frac{d\Gamma}{ds}(\vec{\xi} = \vec{e}) + \frac{d\Gamma}{ds}(\vec{\xi} = -\vec{e})}. \quad (11)$$

After lengthy and straightforward calculations, we get the following expression for $P_T(s)$

$$\begin{aligned} P_T = & -\frac{8\pi m_{\Lambda_b}^3 v \sqrt{s\lambda}}{\left[\mathcal{T}_0(s) + \frac{1}{3}\mathcal{T}_2(s)\right]} \left\{ m_\ell \left(\text{Im}[(A_1 + B_1)^* F_1] + \text{Im}[(A_1 - B_1)^* H_1] \right) \right. \\ & + 8m_{\Lambda_b} \left[2m_\ell \text{Im}[E_1^* C_T f_T^V] - (1 - \sqrt{r}) \left(\text{Im}[H_1^* C_{TE} f_T] + 2m_\ell \text{Im}[E_1^* C_{TE} f_T^V] \right) \right] \\ & - m_\ell m_{\Lambda_b} \left[- (1 - \sqrt{r}) \text{Im}[(A_2 - B_2)^* H_1] + (1 + \sqrt{r}) \text{Im}[(A_2 + B_2)^* F_1] \right] \\ & + 8m_\ell m_{\Lambda_b} \left[(1 - \sqrt{r}) \text{Im}[(D_3 - E_3)^* C_T f_T] + 2(1 + \sqrt{r}) \text{Im}[(D_3 + E_3)^* C_{TE} f_T] \right] \\ & - 16m_\ell m_{\Lambda_b} \left(\text{Im}[D_1^* (C_T + C_{TE}) f_T^V] - \sqrt{r} \text{Im}[D_1^* C_{TE} f_T^V] \right) \\ & + 8m_\ell \left[\text{Im}[(D_1 - E_1)^* C_T f_T] - 2 \text{Im}[(D_1 + E_1)^* C_{TE} f_T] \right] \\ & - m_{\Lambda_b}^3 (1 - \sqrt{r}) (1 + 2\sqrt{r} + r - s) \text{Im}[(D_2 - E_2)^* C_T f_T^S] \\ & + m_{\Lambda_b}^2 (1 - r + s) \left(\text{Im}[A_2^* D_1 - A_1^* D_2] - \text{Im}[B_2^* E_1 - B_1^* E_2] \right) \\ & + 16m_\ell m_{\Lambda_b} \left(\text{Im}[(D_2 - E_2)^* C_T f_T] + 2 \text{Im}[(D_2 + E_2)^* C_{TE} f_T] \right) \\ & - m_{\Lambda_b} s \text{Im}[(D_3 + E_3)^* C_{TE} f_T^V] \\ & + 4m_{\Lambda_b} \left[(1 - \sqrt{r}) \text{Im}[H_2^* C_T f_T] - (1 + \sqrt{r}) \left(\text{Im}[F_1^* C_T f_T] - 2 \text{Im}[F_2^* C_{TE} f_T] \right) \right] \\ & + m_{\Lambda_b} s \left(\text{Im}[F_1 C_T f_T^V] - 2 \text{Im}[F_2 C_{TE} f_T^V] \right) \\ & + 2m_{\Lambda_b} \left(\text{Im}[A_1^* E_1 - B_1^* D_1] - m_{\Lambda_b}^2 s \text{Im}[A_2^* E_2 - B_2^* D_2] \right) \\ & \left. + 8m_\ell m_{\Lambda_b}^2 \left([s - (1 + \sqrt{r})^2] \text{Im}[(D_1 - E_1)^* C_T f_T^S] - (1 - r + s) \text{Im}[(D_2 - E_2)^* C_T f_T^V] \right) \right\}. \quad (12) \end{aligned}$$

It follows from Eq. (12) that, for P_T to have nonvanishing value,

- interactions of new type must exist, and,
- combinations of different Wilson coefficients must have a weak phase.

It should be noted that in the SM $P_T \sim m_s \text{Im}(C_7 C_{10}^*)$ (see [7]) and hence it is strongly suppressed. For this reason, if a large value for P_T is measured in the experiments, it could be considered as an unambiguous indication of the existence of new physics beyond the SM.

It should be remembered that the electromagnetic interaction of final particles can induce P_T , but this contribution is negligibly small, which is of the order of $\mathcal{O}(10^{-3})$.

3 Numerical analysis

In this section we will investigate the sensitivity of the transversal lepton polarization to the new Wilson coefficients. The main input parameters in Eq. (12) are the transition form factors. For transition form factors we will use the results from QCD sum rules approach in combination with HQET [15], which reduces the number of form factors into two independent ones. The dependence of these form factors on s can be expressed as

$$F(q^2) = \frac{F(0)}{1 - a_F s + b_F s^2}, \quad (13)$$

where parameters $F_i(0)$, a and b are listed in table 1.

	$F(0)$	a_F	b_F
F_1	0.462	-0.0182	-0.000176
F_2	-0.077	-0.0685	0.00146

Table 1: Parameters for the form factors given in Eq. (13) in the QCD sum rules method.

In further numerical analysis, for the values of the Wilson coefficients C_7 , C_9^{eff} , and C_{10} we will use next-to leading order logarithmic approximation results at renormalization point $\mu = m_b$ [16]. It should be noted that, in addition to the short distance contribution, C_9^{eff} also receives long distance contributions from the real $\bar{c}c$ resonant states of the J/ψ family. In the present work we do not take into account the long distance effects. Furthermore in carrying out numerical calculations we vary all new Wilson coefficients in the range $0 \leq C_X \leq |C_{10}|$. The experimental bounds on the branching ratio of the $B \rightarrow K^* \mu^+ \mu^-$ and $B \rightarrow \mu^+ \mu^-$ [17] suggest that this is the right order of magnitude range for the vector and scalar interaction coefficients. Moreover we assume that tensor interactions also vary in this region. As we have already noted, in order to obtain considerably large value for P_T , the new Wilson coefficients must have weak phase. For simplicity we assume that all new Wilson coefficients have a common weak phase ϕ .

It follows from the explicit expression of P_T that it depends on s , magnitude and phase of the new Wilson coefficients. The dependence of P_T is eliminated by performing integration over s , i.e., we analyze the averaged P_T , which is defined as

$$\langle P_T \rangle = \frac{\int_{4m_\ell^2/m_{\Lambda_b}^2}^{(1-m_\Lambda/m_{\Lambda_b})^2} P_T \frac{d\mathcal{B}}{ds} ds}{\int_{4m_\ell^2/m_{\Lambda_b}^2}^{(1-m_\Lambda/m_{\Lambda_b})^2} \frac{d\mathcal{B}}{ds} ds}. \quad (14)$$

The dependence of the averaged $\langle P_T \rangle$ on the Wilson coefficients C_T , C_{TE} , C_{RL} , C_{RR} , C_{RLRL} and C_{RLLR} and on the phase ϕ for the $\Lambda_b \rightarrow \Lambda \tau^+ \tau^-$ decay is presented in Figs. (1)–(6). The dependence of $\langle P_T \rangle$ on C_{RL} and on the weak phase for the $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ decay is depicted in Fig. (7). On the other hand, the dependence of $\langle P_T \rangle$ on the remaining Wilson coefficients C_{LL} , C_{LR} , C_{LRRL} and C_{LRLR} for the $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ decay is not presented since for all these choices $\langle P_T \rangle$ is negligibly small ($< 0.2\%$).

It can easily be seen from these figures that $\langle P_T \rangle$ attains at its largest value for C_{RL} about $\sim 25\%$, for $C_{RR} \sim 4\%$, for $C_{TE} \sim 8\%$, for C_{RLRL} , C_{RLLR} about $\sim 5\%$ and about 2% for C_T for the $\Lambda_b \rightarrow \Lambda \tau^+ \tau^-$ decay.

The situation for the $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ decay is similar to the previous case. It is observed that $\langle P_T \rangle$ displays even stronger dependence only on C_{RL} and reaches its maximum value of $\sim 40\%$, and for C_T about 1% , while for all remaining Wilson coefficients other than these two $\langle P_T \rangle$ is negligibly small ($\lesssim 0.2\%$).

These behaviors can be explained as follows. In the massless lepton limit (neglecting scalar type interactions for simplicity) the expression for P_T reduces to

$$P_T \sim \frac{1}{\mathcal{T}_0 + \frac{1}{3}\mathcal{T}_2} \left\{ A \text{Im}[C_{BR}^* (C_{RR} - C_{RL}) - C_{SL}^* (2C_{10} + C_{LR} - C_{LL})] \right. \\ \left. + B \text{Im}[(C_{RL}^* + C_{RR}^*) (2C_{10} + C_{LR} - C_{LL}) + (2C_9 + C_{LR} + C_{LL}) (C_{RR}^* - C_{RL}^*)] \right\} \quad (15)$$

where A and B are functions of s and ratio of the form factors. In this work we will assume that C_{BR} and C_{SL} are defined in the same way as in the SM, since the measured branching ratio of $B \rightarrow X_s \gamma$ decay is in a very good agreement with the SM prediction [18]–[20]. The first term in Eq. (15) can safely be neglected since $|C_7^{SM}| \ll |C_9^{eff}|$ and $|C_7^{SM}| \ll |C_{10}|$. We also observe that the expression in Eq. (15) involves C_{RR} and C_{RL} in the combinations of the form

$$\sim \text{Im}[C_{RR}^* (C_{10} + C_9^{eff})] , \\ \sim \text{Im}[C_{RL}^* (C_{10} - C_9^{eff})] .$$

In the SM C_{10} and C_9^{eff} have opposite signs and have almost equal magnitudes (see for example [21]), which explains the reason why P_T gets larger values for C_{RL} compared to the other new Wilson coefficients. For all other choices P_T is proportional to the lepton mass and for this reason P_T is quite small the coefficients C_{LL} , C_{LR} , C_{LRRL} and C_{LRLR} .

Finally we would like to discuss briefly the detectability of $\langle P_T \rangle$ in the experiments. Experimentally, to measure $\langle P_T \rangle$ of a particular decay with the branching ratio $\mathcal{B}$ at the $n\sigma$ level, the required number of events are $N = n^2/(\mathcal{B} \langle P_T \rangle)^2$. For example, if $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-) \sim 10^{-6}$ then, to measure $\langle P_T \rangle \simeq 0.2$ at 3σ level, $N \simeq 4.5 \times 10^7$ Λ_b decays are required. Since at LHC and BTeV machines 10^{12} $b\bar{b}$ pairs are expected to be produced per year [?], the detectability of $\langle P_T \rangle$ is quite high in these colliders.

References

- [1] CPLEAR Collaboration, A. Angelopoulos *et. al* *Phys. Lett.* **B444** (1998) 43.
- [2] N. Cabibbo, *Phys. Rev. Lett.* **10** (1963) 531;
M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49** (1973) 652.
- [3] C. Caso *et. al*, *Eur. J. Phys.* **15** (2000) 1.
- [4] The BaBar Physics book, Eds: P. F. Harrison, H. R. Quinn, SLAC report (1998) 504;
BELLE Collaboration, E. Drebyš *et. al*, *Nucl. Inst. MEthods* **A446** (2000) 89.
- [5] B. Aubert *et. al*, BaBar Collaboration, hep-ex/0203007;
K. Trabelsi, DELLE Collaboration, talk given at the XXXVII th Recontres de Moriond
Electroweak Interactions and Unified Theories, 2002.
- [6] Chuan-Hung Chen, C. Q. Geng and J. N. Ng, *Phys. Rev.* **D65** (2002) 091502.
- [7] Chuan-Hung Chen, C. Q. Geng, preprint: hep-ph/0205306 (2002).
- [8] T. M. Aliev, A. Özpineci, M. Savcı, *Phys. Rev.* **D** (in press).
- [9] T. M. Aliev, M. K. Çakmak, M. Savcı, *Nucl. Phys.* **B607** (2001) 305.
- [10] T. M. Aliev, C. S. Kim, Y. G. Kim, *Phys. Rev.* **D62** (2000) 014026.
- [11] S. Fukae, C. S. Kim, T. Yoshikawa, *Phys. Rev.* **D61** (2000) 074015.
- [12] T. M. Aliev, A. Özpineci, M. Savcı, preprint: hep-ph/0202120 (2002).
- [13] T. Mannel, W. Roberts and Z. Ryzak, *Nucl. Phys.* **B355** (1991) 38.
- [14] Chuan-Hung Chen and C. Q. Geng, *Phys. Rev.* **D63** (2001) 054005; *Phys. Rev.* **D63**
(2001) 114024; *Phys. Rev.* **D64** (2001) 074001.
- [15] C.-S. Huang, H.-G. Yan, *Phys. Rev.* **D59** (1999) 114022.
- [16] C. Bobeth, M. Misiak, and J. Urban, *Nucl. Phys.* **B574** (2000) 291.
- [17] CDF Collaboration, T. Affolder *at. al*, *Phys. Rev. Lett.* **83** (1999) 3378;
CLEO Collaboration, Yongsheng Gao, prep. hep-ex/0108005 (2001).
- [18] M. S. Alam *et. al*, CLEO Collaboration, *Phys. Rev. Lett.* **74** (1995) 2885;
S. Ahmed *at. al*, CLEO Collaboration, hep-ex/9908022;
D. Cassel, CLEO Collaboration, talk at Lepton Photon Conf. 2001, Rome, July 2001.
- [19] R. Barate *at. al*, ALEPH Collaboration, *Phys. Lett.* **B429** (1998) 169.
- [20] K. Abe *at. al*, BELLE Collaboration, *Phys. Lett.* **B511** (2001) 151.
- [21] G. Buchalla, A. J. Buras and M. E. Lautenbacher, *Rev. Mod. Phys.* **68** (1996) 1230.
- [22] N. Harnew, proceedings of heavy flavors 8, (Southampton, 1999).

Figure captions

Fig. (1) The dependence of the averaged transversal lepton polarization $\langle P_T^- \rangle$ on the phase ϕ and on the new Wilson coefficient C_T for the $\Lambda_b \rightarrow \Lambda \tau^+ \tau^-$ decay.

Fig. (2) The same as in Fig. (1), but for the Wilson coefficient C_{TE} .

Fig. (3) The same as in Fig. (1), but for the Wilson coefficient C_{RL} .

Fig. (4) The same as in Fig. (1), but for the Wilson coefficient C_{RR} .

Fig. (5) The same as in Fig. (1), but for the Wilson coefficient C_{RLRL} .

Fig. (6) The same as in Fig. (1), but for the Wilson coefficient C_{RLLR} .

Fig. (7) The dependence of the averaged transversal lepton polarization $\langle P_T^- \rangle$ on the phase ϕ and on the new Wilson coefficient C_{RL} for the $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$ decay.




Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:




Figure 6:




Figure 7: